



# Refined Neutrosophic Matrices Representations

Mohammad Abobala

## ► To cite this version:

| Mohammad Abobala. Refined Neutrosophic Matrices Representations. 2021. hal-03265163

**HAL Id: hal-03265163**

**<https://hal.science/hal-03265163>**

Preprint submitted on 19 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Refined Neutrosophic Matrices Representations

Mohammad Abobala,

Tishreen University, Faculty of Science, Department of Mathematics, Latakia, Syria

## Abstract:

This paper studies the possibility of representing refined neutrosophic matrices by linear functions over a refined neutrosophic vector space.

## Main Discussion

### Definition 1:

Let  $V, W$  be two vector spaces over  $R$ , consider any three linear transformations  $g, h, q: V \rightarrow W$ . We define the corresponding full AH-refined linear transformation as follows:

$$f: V(I_1, I_2) \rightarrow W(I_1, I_2),$$

$$f(x + yI_1 + zI_2) = g(x) + I_1[(g + h + q)(x + y + z) - (g + q)(x + z)] + I_2[(g + q)(x + z) - g(x)].$$

We denote it by  $f = g + hI_1 + qI_2$

### Theorem 2:

Let  $f = g + hI_1 + qI_2: V(I_1, I_2) \rightarrow W(I_1, I_2)$  be any full AH-refined linear transformation, then  $f$  is linear by classical meaning.

### Proof:

$\forall X = x + yI_1 + zI_2, Y = x_1 + y_1I_1 + z_1I_2 \in V(I_1, I_2)$  we have:

$$\begin{aligned} f(X + Y) &= f[(x + x_1) + I_1(y + y_1) + I_2(z + z_1)] \\ &= g(x + x_1) + I_1[(g + h + q)(x + x_1 + y + y_1 + z + z_1) - (g + q)(x + x_1 + z + z_1)] \\ &\quad + I_2[(g + q)(x + x_1 + z + z_1) - g(x + x_1)] \\ &= g(x) + g(x_1) \\ &\quad + I_1[(g + h + q)(x + y + z) + (g + h + q)(x_1 + y_1 + z_1) - (g + q)(x + z) \\ &\quad - (g + q)(x_1 + z_1)] + I_2[(g + q)(x + z) + (g + q)(x_1 + z_1) - g(x) - g(x_1)] \\ &= [g(x) + I_1[(g + h + q)(x + y + z) - (g + q)(x + z)] + I_2[(g + q)(x + z) - g(x)]] + [g(x_1) + \\ &\quad I_1[(g + h + q)(x_1 + y_1 + z_1) - (g + q)(x_1 + z_1)] + I_2[(g + h + q)(x_1 + z_1) - g(x_1)]], \\ &= f(X) + f(Y) \end{aligned}$$

$\forall r = r_0 + r_1I_1 + r_2I_2 \in V(I_1, I_2)$ , we have:

$$r \cdot X = r_0x + I_1[(r_0 + r_1 + r_2)(x + y + z) - (r_0 + r_2)(x + z)] + I_2[(r_0 + r_2)(x + z) - r_0x]$$

$$f(r \cdot X) = g(r_0x) + I_1[(g + h + q)(r_0 + r_1 + r_2)(x + y + z) - (g + q)(r_0 + r_2)(x + z)] + I_2[(g + q)(r_0 + r_2)(x + z) - g(r_0x)],$$

Now we compute  $r \cdot f(X)$ :

$$\begin{aligned} r \cdot f(X) &= (r_0 + r_1I_1 + r_2I_2) \cdot [g(x) + I_1[(g + h + q)(x + y + z) - (g + q)(x + z)] + I_2[(g + q)(x + z) - g(x)], \\ &= r_0 g(x) + I_1[(r_0 + r_1 + r_2)(g + h + q)(x + y + z) - (r_0 + r_2)(g + q)(x + z)] + I_2[(r_0 + r_2) \cdot (g + q)(x + z) - r_0 g(x)] = f(rX). \end{aligned}$$

**Definition 3:**

Let  $M = A + BI_1 + CI_2$  be a matrix over  $R(I_1, I_2)$ . We say that  $M$  is the matrix of  $f = g + hI_1 + qI_2$  iff  $f(x + yI_1 + zI_2) = M \cdot (x + yI_1 + zI_2)$ .

**Theorem 4:**

Let  $f = g + hI_1 + qI_2: V(I_1, I_2) \rightarrow W(I_1, I_2)$  be any full AH-refined linear transformation, then  $M = A + BI_1 + CI_2$  is the matrix of  $f$  iff  $A$  is the matrix of  $g$ ,  $B$  is the matrix of  $h$ ,  $C$  is the matrix of  $q$ .

**Proof:**

Firstly, we suppose that  $A$  is the matrix of  $g$ ,  $B$  is the matrix of  $h$ ,  $C$  is the matrix of  $q$ , where  $g, h, q: V \rightarrow W$  are classical linear transformations.

Consider the refined neutrosophic matrices  $M = A + BI_1 + CI_2$ , we have for each  $x + yI_1 + zI_2 \in V(I_1, I_2)$ :

$$\begin{aligned} M(x + yI_1 + zI_2) &= (A + BI_1 + CI_2)(x + yI_1 + zI_2) = A \cdot x + I_1[(A + B + C)(x + y + z) - (A + C)(x + z)] + I_2[(A + C)(x + z) - A \cdot x], \\ &= g(x) + I_1[(g + h + q)(x + y + z) - (g + q)(x + z)] + I_2[(g + q)(x + z) - g(x)] = f(x + yI_1 + zI_2), \text{ hence } M \text{ is the matrix of } f. \end{aligned}$$

Conversely, we assume that  $M = A + BI_1 + CI_2$  is the matrix of  $f$ , we must prove that  $A$  is the matrix of  $g$ ,  $B$  is the matrix of  $h$ ,  $C$  is the matrix of  $q$ .

$$\begin{aligned} \text{We have } f(x + yI_1 + zI_2) &= M(x + yI_1 + zI_2), \text{ thus: } g(x) + I_1[(g + h + q)(x + y + z) - (g + q)(x + z)] + I_2[(g + q)(x + z) - g(x)] \\ &= Ax + I_1[(A + B + C)(x + y + z) - (A + C)(x + z)] + I_2[(A + C)(x + z) - Ax] \end{aligned}$$

$$\text{So that, } g(x) = Ax, (A + C)(x + z) = (g + q)(x + z), (A + B + C)(x + y + z) = (g + h + q)(x + y + z)$$

This implies that  $A$  is the matrix of  $g$ ,  $A + C$  is the matrix of  $g + q$ , thus  $C$  is the matrix of  $q$  and  $A + B + C$  is the matrix of  $g + h + q$ , thus  $B$  is the matrix of  $h$ .

In order to prove that every refined neutrosophic matrix can be represented by a unique AH-refined linear transformation, we introduce the following algorithm to derive a basis of  $V(I_1, I_2)$  from any classical basis of  $V$ .

**Theorem 5:** Let  $S = \{V_1, V_2, \dots, V_n\}$  be a basis of  $V$  over  $R$ , then:

$$L_{i,j,k} = \{V_i + (V_j - V_k)I_1 + (V_k - V_i)I_2\}$$

Is a basis of  $V(I_1, I_2)$  over  $R(I_1, I_2)$ .

**Theorem 6:** Let  $f: V(I_1, I_2) \rightarrow W(I_1, I_2)$  be a linear function, then  $f$  must be a full AH-refined linear transformation.

**Example 7:**

Now, we give an example to clarify the structure of the refined neutrosophic basis of a refined neutrosophic vector space basing on Theorem 5.

It is well known that the  $R^2$  is a vector space over the real field  $R$ . Consider the corresponding refined neutrosophic vector space

$$R^2(I_1, I_2) = \{a + bI_1 + cI_2; a, b, c \in R^2\} = \{(x, y) + I_1(z, t) + I_2(m, n); m, n, z, t, x, y \in R\}.$$

The basis of  $R^2$  is the set  $\{e_1 = (1, 0), e_2 = (0, 1)\}$ . According to Theorem 5, the related refined neutrosophic basis of  $R^2(I_1, I_2)$  is:

$$v_1 = e_1 + I_1(e_1 - e_1) + I_2(e_1 - e_1), v_2 = e_1 + I_1(e_2 - e_1) + I_2(e_1 - e_1)$$

$$v_3 = e_1 + I_1(e_2 - e_2) + I_2(e_2 - e_1), v_4 = e_1 + I_1(e_1 - e_2) + I_2(e_2 - e_1),$$

$$v_5 = e_2 + I_1(e_1 - e_1) + I_2(e_1 - e_2), v_6 = e_2 + I_1(e_1 - e_2) + I_2(e_2 - e_2),$$

$$v_7 = e_2 + I_1(e_2 - e_1) + I_2(e_1 - e_2), v_8 = e_2 + I_1(e_2 - e_2) + I_2(e_2 - e_2).$$

**Example 8:**

Now, we illustrate an example to clarify the representation of refined linear transformations by refined neutrosophic matrices.

$$\text{Let } f: R^2(I_1, I_2) \rightarrow R(I_1, I_2); f[(x, y) + I_1(z, t) + I_2(m, n)] = f[(x + zI_1 + mI_2, y + tI_1 + nI_2)] = \\ 2(x + zI_1 + mI_2) - (y + tI_1 + nI_2) = (2x - y) + (2z - t)I_1 + (2m - n)I_2.$$

be a refined linear transformation, we have to write  $f$  as an AH-refined linear transformation.

By Theorem 6, we know that  $f$  must be equal to  $g + hI_1 + qI_2$ , where  $g, h, q: R^2 \rightarrow R$ .

Firstly, we have  $g: R^2 \rightarrow R; g(x, y) = 2x - y$ , and  $(g + q)[(x, y) + (m, n)] - g(x, y) = 2m - n$ ,

This means that:  $g(x + m, y + n) + q(x + m, y + n) - (2x - y) = 2m - n$ , so that

$$2(x + m) - (y + n) + q(x + m, y + n) - (2x - y) = 2m - n, \text{ hence}$$

$$2m - n + q(x + m, y + n) = 2m - n, \text{ thus } q(x + m, y + n) = 0 \text{ for all } x, y, m, n.$$

This implies that  $q(x, y) = 0$  is a zero map.

For the computing of  $h$ , we regard that:  $(g + h + q)[(x, y) + (z, t) + (m, n)] - (g + q)[(x, y) + (m, n)] = 2z - t$ .

So that:  $(g + q + h)[(x + z + m, y + t + n)] - (g + q)[(x + m, y + n)] = 2z - t$ . By remarking that

$g + q = g$ , we get:

$$(g + h)[(x + z + m, y + t + n)] - g[(x + m, y + n)] = 2z - t, \text{ hence}$$

$$g(x + z + m, y + t + n) + h(x + z + m, y + t + n) - [2(x + m) - (y + n)] = 2z - t,$$

$$\text{This means that: } 2(x + z + m) - (y + t + n) + h(x + z + m, y + t + n) - 2x - 2m + y + n = 2z - t,$$

Thus  $h(x + z + m, y + t + n) = 0$  for all  $x, y, z, m, n, t$ , i.e.  $h$  is a zero map.

By the previous discussion, we get that  $f = g + 0.I_1 + 0.I_2$ .

The matrix of  $g$  is  $\begin{pmatrix} 2 \\ -1 \end{pmatrix}$ , the matrix of  $q$  and  $h$  is  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ , thus the refined neutrosophic matrix of  $f$  is

$$\begin{pmatrix} 2 \\ -1 \end{pmatrix} + I_1 \begin{pmatrix} 0 \\ 0 \end{pmatrix} + I_2 \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix} = M.$$

Now, we check that  $M$  is refined neutrosophic matrix of  $f$  by the following computing:

$$M. [(x, y) + I_1(z, t) + I_2(m, n)] = M. [(x + zI_1 + mI_2, y + tI_1 + nI_2)] = 2(x + zI_1 + mI_2) - (y + tI_1 + nI_2) = f(x + zI_1 + mI_2, y + tI_1 + nI_2).$$

#### Example 9:

In example 8, we showed how a refined neutrosophic linear transformation can be turned into a refined neutrosophic matrix, now we do the converse. We illustrate an example to show how a refined neutrosophic matrix can be turned into a refined neutrosophic linear transformation.

Let  $M = \begin{pmatrix} 1 & I_2 \\ 3 - I_1 + 2I_2 & 2I_1 \end{pmatrix}$  be a refined neutrosophic matrix.  $M$  can be written in the following form:

$$M = A + BI_1 + CI_2 = \begin{pmatrix} 1 & 0 \\ 3 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ -1 & 2 \end{pmatrix} I_1 + \begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix} I_2.$$

$M$  can be represented by a unique refined neutrosophic linear transformation

$$f: R^2(I_1, I_2) \rightarrow R^2(I_1, I_2); f[(x + zI_1 + mI_2, y + tI_1 + nI_2)] = M. [(x + zI_1 + mI_2, y + tI_1 + nI_2)] = (x + I_1(z + t) + I_2(m + y + n), 3x + (4z - x + 2y + 2t + 2n - m)I_1 + (2x + 5m)I_2).$$

#### References

- [1] Abobala, M., "On Some Special Substructures of Neutrosophic Rings and Their Properties", International Journal of Neutrosophic Science", Vol. 4, pp. 72-81, 2020.
- [2] Abobala, M., "On Some Special Substructures of Refined Neutrosophic Rings", International Journal of Neutrosophic Science, Vol. 5, pp. 59-66, 2020.

- [3] Abobala, M., "Classical Homomorphisms Between Refined Neutrosophic Rings and Neutrosophic Rings", International Journal of Neutrosophic Science, Vol. 5, pp. 72-75, 2020.
- [4] Abobala, M., and Alhamido, R., "AH-Substructures in Neutrosophic Modules", International Journal of Neutrosophic Science, Vol. 7, pp. 79-86 , 2020.
- [5] Agboola, A.A.A., "On Refined Neutrosophic Algebraic Structures", Neutrosophic Sets and Systems, Vol.10, pp. 99-101, 2015.
- [6] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol. 2(2), pp. 77-81, 2020.
- [7] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings II", International Journal of Neutrosophic Science, Vol. 2(2), pp. 89-94, 2020.
- [8] Kandasamy, V. W. B., Ilanthenral, K., and Smarandache, F., " Semi-Idempotents in Neutrosophic Rings", Mathematics Journal (MDPI), Vol. 7, 2019.
- [9] Abobala, M., "AH-Subspaces in Neutrosophic Vector Spaces", International Journal of Neutrosophic Science, Vol. 6 , pp. 80-86, 2020.
- [10] Smarandache, F., " A Unifying Field in Logics: Neutrosophic Logic, Neutrosophy, Neutrosophic Set., Neutrosophic Probability", American Research Press, Rehoboth, 2003.
- [11] Sankari, H., and Abobala, M., "Solving Three Conjectures About Neutrosophic Quadruple Vector Spaces", Neutrosophic Sets and Systems, Vol. , pp. , 2020.
- [12] Smarandache, F., and Abobala, M., "n-Refined neutrosophic Rings", International Journal of Neutrosophic Science, Vol. , pp. , 2020.
- [13]. Sankari, H., and Abobala, M., "n-Refined Neutrosophic Modules", Neutrosophic Sets and Systems, Vol.36, pp. 1-11, 2020.
- [14] Sankari, H., and Abobala, M., "AH-Homomorphisms In Neutrosophic Rings and Refined Neutrosophic Rings", Neutrosophic Sets and Systems, 2020.
- [15]. Abobala, M., "n- Refined Neutrosophic Groups II", International journal Of Neutrosophic Science, Vol. 0, 2019.

- [16]. Abobala, M., " A Review On Recent Developments of Neutrosophic Matrix Theory and Open Problems", hal- Archives, 2021.
- [17]. Abobala, M., Partial Foundation of Neutrosophic Number Theory, Neutrosophic Sets and Systems, Vol. 39 , 2021
- [18]. Abobala, M., " Semi Homomorphisms and Algebraic Relations Between Strong Refined Neutrosophic Modules and Strong Neutrosophic Modules", Neutrosophic Sets and Systems, Vol. 39, 2021.
- [19] Abobala, M., "A Study of AH-Substructures in  $n$ -Refined Neutrosophic Vector Spaces", International Journal of Neutrosophic Science", Vol. 9, pp.74-85. 2020.
- [20]. Abobala, M., "Maximal and Minimal Ideals In Some Neutrosophic Rings", Hal-Archives, 2021.
- [21] Aswad. F, M., " A Study of neutrosophic Bi Matrix", Neutrosophic Knowledge, Vol. 2, 2021
- [22] Hatip, A., and Abobala, M., "AH-Substructures In Strong Refined Neutrosophic Modules", International Journal of Neutrosophic Science, Vol. 9, pp. 110-116 . 2020.
- [23] Abobala, M., " Classical Homomorphisms Between  $n$ -refined Neutrosophic Rings", International Journal of Neutrosophic Science", Vol. 7, pp. 74-78 , 2020.
- [24] Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona 2006.
- [25] Smarandache F., and Abobala, M., "  $n$ -Refined Neutrosophic Vector Spaces", International Journal of Neutrosophic Science, Vol. 7, pp. 47-54, 2020.
- [26] Hatip, A., Alhamido, R., and Abobala, M., " A Contribution to Neutrosophic Groups", International Journal of Neutrosophic Science", Vol. 0, pp. 67-76 , 2019.
- [27] Abobala, M., "On The Representation of Neutrosophic Matrices by Neutrosophic Linear Transformations", Journal of Mathematics, Hindawi, 2021
- [28] Abobala, M., "On Some Algebraic Properties of  $n$ -Refined Neutrosophic Elements and  $n$ -Refined Neutrosophic Linear Equations", Mathematical Problems in Engineering, Hindawi, 2021
- [29] Abobala, M., Hatip, A., Olgun, N., Broumi, S., Salama, A.A., and Khaled, E, H., "The Algebraic Creativity In The Neutrosophic Square Matrices", Neutrosophic Sets and Systems, Vol. 40, pp. 1-11, 2021.
- [30] Abobala, M., "Neutrosophic Real Inner Product Spaces", Neutrosophic Sets and Systems, vol. 43, 2021.

- [31] Abobala, M., "A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings", International Journal of Mathematics and Mathematical Sciences, hindawi, 2021
- [32] Abobala, M., and Hatip, A., "An Algebraic Approach to Neutrosophic Euclidean Geometry", Neutrosophic Sets and Systems, Vol. 43, 2021.
- [33] Abobala, M., "On Some Neutrosophic Algebraic Equations", Journal of New Theory, Vol. 33, 2020.
- [34] Abobala, M., " $n$ -Cyclic Refined Neutrosophic Algebraic Systems Of Sub-Indeterminacies, An Application To Rings and Modules", International Journal of Neutrosophic Science, Vol. 12, pp. 81-95 . 2020.
- [35] Abobala, M., "On Some Special Elements In Neutrosophic Rings and Refined Neutrosophic Rings", Journal of New Theory, vol. 33, 2020.
- [36] Abobala, M., "  $n$ -Refined Neutrosophic Groups I", International Journal of Neutrosophic Science, Vol. 0, pp. 27-34. 2020
- [37] Sankari, H., and Abobala, M., "Neutrosophic Linear Diophantine Equations With two Variables", Neutrosophic Sets and Systems, Vol. 38, pp. 22-30, 2020.
- [38] Abobala, M., " A Study Of Maximal and Minimal Ideals Of  $n$ -Refined Neutrosophic Rings", Journal of Fuzzy Extension and Applications, 2021.