



## Synthesis and characterization of group 6-9 metal-rich homo- and hetero-metallaboranes

Ranjit Bag, Suman Gomosta, Alaka Nanda Pradhan, Thierry Roisnel,  
Sundargopal Ghosh

### ► To cite this version:

Ranjit Bag, Suman Gomosta, Alaka Nanda Pradhan, Thierry Roisnel, Sundargopal Ghosh. Synthesis and characterization of group 6-9 metal-rich homo- and hetero-metallaboranes. *Journal of the Indian Chemical Society*, 2021, 98 (3), pp.100040. 10.1016/j.jics.2021.100040 . hal-03269459


**HAL Id: hal-03269459**

**<https://univ-rennes.hal.science/hal-03269459>**

Submitted on 30 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



J. Indian Chem. Soc.

ISSN: 0019-4522

Vol..../Issue.../pp...



## Synthesis and characterization of group 6-9 metal-rich homo- and hetero-metallaboranes

Ranjit Bag,<sup>a</sup> Suman Gomosta,<sup>a</sup> Alaka Nanda Pradhan,<sup>a</sup> Thierry Roisnel,<sup>b</sup> and Sundargopal Ghosh<sup>a\*</sup>

<sup>a</sup>Department of Chemistry, Indian Institute of Technology Madras, Chennai 600036, India. Tel. +91 44-22574230; Fax. +91 44-22574202.

<sup>b</sup>Université de Rennes, CNRS, Institut des Sciences Chimiques de Rennes, UMR 6226, F-35000 Rennes, France.

\*E-mail: sghosh@iitm.ac.in

Manuscript Received online: ..... (Date), Accepted: ... (Date), Published online ..... (Date)

**Abstract:** To isolate the metal-rich metallaboranes of group 6-9, we have performed the reaction of various reaction intermediates, generally synthesized from the low-temperature reactions of  $[\text{Cp}^*\text{WCl}_4]$  ( $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$ ),  $[(\text{Cp}^*\text{RhCl}_2)_2]$ , or  $[(\text{Cp}^*\text{RuCl}_2)_2]$  and  $[\text{LiBH}_4 \cdot \text{THF}]$  with different transition metal carbonyl compounds. For example, the thermolytic reaction of  $[\text{Fe}_2(\text{CO})_9]$  with an *in situ* generated intermediate, produced from the reaction of  $[\text{Cp}^*\text{WCl}_4]$  and  $[\text{LiBH}_4 \cdot \text{THF}]$  afforded a trigonal bipyramidal cluster,  $[(\mu_3\text{-BH})_2\text{H}_2\{\text{Cp}^*\text{W}(\text{CO})_2\}\{\text{Cp}^*\text{W}(\text{CO})\}\{\text{Fe}(\text{CO})_3\}]$ , **1** which contains a triply-bridging *bis*-{hydrido(borylene)} ligand. Similarly, the reaction of  $[\text{Co}_2(\text{CO})_8]$  with *nido*- $[(\text{RhCp}^*)_2(\text{B}_3\text{H}_7)]$  **I** at room temperature, yielded an octahedral cluster,  $[(\text{Cp}^*\text{Rh})_2\text{B}_2\text{H}_2\text{Co}_2(\text{CO})_5(\mu_3\text{-CO})]$ , **2**. In this reaction, *nido*-**I** having (n+2) skeletal electron pairs (SEP) goes on for the formation of a *closo*-rhodaborane with (n+1) SEP. In addition, we have isolated a trinuclear bis( $\mu_3$ -oxo) metalla cluster  $[(\text{Cp}^*\text{Ru})_3(\mu_3\text{-OBF}_3)_2(\mu\text{-H})]$ , **3**. Compound **3** can be considered as cluster having trigonal bipyramidal geometry with *exo*- $\text{BF}_3$  fragment. All these clusters were characterized by IR, mass spectrometry, NMR, and single-crystal X-ray crystallographic analysis.



**Keywords:** borylene, tungsten, metal-rich metallaborane, octahedral complex, oxoruthena cluster


## Introduction

The chemistry of metallaborane cluster has developed rapidly in the last few decades due to their diversity of fascinating structures, unique bonding and reactions.<sup>1-3</sup> Several approaches are there for the cluster networks expansion,<sup>4-5</sup> among them two major approaches are metathesis reaction with monoborane reagents<sup>6-7</sup> and insertion of metal carbonyl fragments into unsaturated metallaboranes.<sup>8</sup> The former method is highly effective for the production of metallaborane compounds with higher vertex.<sup>9-10</sup> However, the latter process is more vibrant as it can act as a source of both metal carbonyl fragment or CO to form a heterometallic metallaborane clusters.<sup>11</sup>

Based on the metal to boron ratio, metallaborane compounds are classified into two main areas boron-rich<sup>12-14</sup> and metal-rich metallaboranes.<sup>15-16</sup> Unfortunately, the great majority of metallaboranes in most of

the synthetic methods have low metal to boron ratios. However, significant growth of metal-rich metallaboranes has been demonstrated by several reviews and articles in recent years.<sup>17-18</sup> Starting from borylenes, multicentred borides and higher vertex clusters, metal-rich metallaboranes have been synthesized in various structural forms. The synthesis of metal-rich metallaboranes are of specific interest owing to their several applications as super-hard materials<sup>19</sup> superconductors<sup>20</sup> and magnetic materials.<sup>21</sup>

In the course of our continuing interest to synthesize various metallaborane clusters, we have been exploring numerous approaches using different metal carbonyl compounds.<sup>22</sup> The reaction of *nido*-metallaboranes with transition metal carbonyl compounds have been found to be the most fruitful means of production of metal-rich metallaboranes.<sup>16a,e</sup>



**Scheme 1.** Examples of some metal-rich metallaboranes.

Thus, we were interested in exploring the reactivity of  $[\text{Cp}^*\text{WCl}_4]$  ( $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$ ),  $(\text{Cp}^*\text{RhCl}_2)_2$ , and  $(\text{Cp}^*\text{RuCl}_2)_2$  with borane reagents and metal carbonyl complexes such as  $\text{Fe}_2(\text{CO})_9$  and  $\text{Co}_2(\text{CO})_8$ . In this report, we have described the synthesis and structure of various metal-rich metallaboranes.

## Experimental

### General Procedures and Instrumentation.

All the experiments were conducted under argon atmosphere using standard Schlenk line and glovebox techniques.

Purification of common Solvents (Hexane, Toluene, THF) were achieved by distillation from suitable drying agents like sodium/benzophenone under argon.  $\text{C}_6\text{D}_6$  and  $\text{CDCl}_3$  solvents were stored over activated molecular sieves.  $[\text{Cp}^*\text{WCl}_4]$ ,<sup>23</sup> *nido*- $[(\text{Cp}^*\text{Rh})_2\text{B}_3\text{H}_7]$ , **I**,<sup>24</sup> and *nido*- $[(\text{Cp}^*\text{Ru})_2\text{B}_3\text{H}_9]$ , **II**<sup>24</sup> were synthesized according the method mentioned in literature.



Chemicals such as,  $[\text{LiBH}_4 \cdot \text{THF}]$  (2.0 M in THF),  $n\text{BuLi}$ ,  $\text{Cp}^*\text{H}$ ,  $[\text{Co}_2(\text{CO})_8]$ , and  $[\text{Fe}_2(\text{CO})_9]$  were obtained from Sigma-Aldrich and used without any further purifications. Iodomethane ( $\text{MeI}$ ) was obtained from Sigma-Aldrich and distilled prior to use. NMR spectra of all the compounds were recorded on Bruker FT-NMR spectrometers (400 or 500 MHz). The residual solvents protons were used as reference ( $\delta$ , ppm,  $\text{CDCl}_3$  – 7.26,  $\text{C}_6\text{D}_6$  – 7.16) and a sealed tube containing  $[\text{Bu}_4\text{N}(\text{B}_3\text{H}_8)]$  ( $\delta_{\text{B}}$ , -30.07 ppm,) in  $\text{C}_6\text{D}_6$  was used as external reference in  $^{11}\text{B}$  NMR, which was synthesized according to the method in the literature.<sup>25</sup> Thin-layer chromatography (TLC) was performed on 250 mm diameter aluminium-supported silica gel plates (Merck 105554 TLC silica gel 60 F<sub>254</sub>). The infrared spectra (IR) were recorded on a FT/IR-4100 JASCO spectrometer in  $\text{CH}_2\text{Cl}_2$  solvent. Electrospray mass (ESI-MS) spectra were recorded on an Agilent 6545 Q-TOF LC/MS instrument.

**Synthesis of 1:** In a flame-dried Schlenk tube  $[\text{Cp}^*\text{WCl}_4]$  (0.1 g, 0.21 mmol) in 10 mL toluene was treated with 5-fold of  $[\text{LiBH}_4 \cdot \text{THF}]$  (0.7 mL) at -78 °C and stir for 1 h in room temperature. The residue was extracted into hexane after removal of toluene and filtered through a frit using celite. The brownish-green hexane solution was dried in vacuo and taken in 10 mL of THF. The solution is heated at 60 °C for 16 h in presence of  $[\text{Fe}_2(\text{CO})_9]$  (0.02 g). The solvent of reaction mixture was evaporated in vacuo and the residue was extracted into hexane, then passed through celite. After the removal of solvent from filtrate, the residue was subjected to chromatographic purification using silica gel TLC plates. Elution with a mixture of hexane:dichloromethane (90:10 v/v) produced yellow solid **1** (0.012 g, 12 %) along with known compounds  $[\{\text{Cp}^*\text{W}(\text{CO})_2\}_2\text{H}_2(\mu_3\text{-BH})_2\{\text{Fe}(\text{CO})_2\}]^{26}$  (0.010 g, 10 %),  $[\{\text{Cp}^*\text{W}(\text{CO})_2\}_2(\mu_3\text{-BH})_2\{\text{Fe}(\text{CO})_3\}]^{26}$  (0.020 g, 20 %), and  $[(\text{Cp}^*\text{W})_2\text{B}_5\text{H}_9]^{27}$  (0.015 g, 20 %).



**1:** MS (ESI<sup>+</sup>):  $m/z$  calculated for [C<sub>26</sub>H<sub>34</sub>B<sub>2</sub>FeO<sub>6</sub>W<sub>2</sub>-(3H+2CO)]<sup>+</sup>: 829.0804, found: 829.0269. <sup>11</sup>B{<sup>1</sup>H} NMR (128MHz, CDCl<sub>3</sub>): 87.9 (s, 1B), 62.4 (s, 1B) ppm. <sup>1</sup>H NMR (500MHz, CDCl<sub>3</sub>): 7.72 (br, 1H, BH<sub>t</sub>), 7.54 (br, 1H, BH<sub>t</sub>), 2.19 (s, 15H, C<sub>5</sub>Me<sub>5</sub>), 1.97 (s, 15H, C<sub>5</sub>Me<sub>5</sub>), -10.64 (1H, W-H-B), -12.44 (1H, W-H-B), ppm. <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): 218.4 (CO), 98.7 (s, C<sub>5</sub>Me<sub>5</sub>), 98.1 (s, C<sub>5</sub>Me<sub>5</sub>), 11.7 (s, C<sub>5</sub>Me<sub>5</sub>), 10.2 (s, C<sub>5</sub>Me<sub>5</sub>) ppm. FTIR (CH<sub>2</sub>Cl<sub>2</sub>): 2415 (BH<sub>t</sub>); 2014, 1943 (CO) cm<sup>-1</sup>.

**Synthesis of 2:** In a flame-dried Schlenk tube, *nido-I* (0.15 g, 0.29 mmol) in 15 mL hexane was reacted with [Co<sub>2</sub>(CO)<sub>8</sub>] (0.198 g, 0.58 mmol) at room temperature and stir for 24 h. The residue was extracted into hexane after removal of solvent and passed through a frit using celite. The mother liquor was kept at -40 °C to remove [Co<sub>4</sub>(CO)<sub>12</sub>] by filtration of mother liquor. After the removal of solvent from filtrate, the residue was subjected to chromatographic purification using silica gel TLC plates. Elution with a

mixture of hexane:dichloromethane (80:20 v/v) in inert atmosphere produced black solid **2** (0.036 g, 16 %) along with a known compound [(Cp\**Rh*)<sub>2</sub>(μ-CO)(μ<sub>3</sub>-BH)<sub>2</sub>{Co(CO)<sub>4</sub>}]<sup>26</sup> (0.044 g, 20%).

**2:** HRMS (ESI<sup>+</sup>):  $m/z$  calculated for [C<sub>26</sub>H<sub>32</sub>O<sub>6</sub>B<sub>2</sub>Co<sub>2</sub>Rh<sub>2</sub>+Na]<sup>+</sup>: 808.9056, found: 808.9052. <sup>11</sup>B{<sup>1</sup>H} NMR (128 MHz, C<sub>6</sub>D<sub>6</sub>): 168.0 (s, 1B), 152.5 (s, 1B) ppm. <sup>1</sup>H NMR (500MHz, C<sub>6</sub>D<sub>6</sub>): 9.35 (br, 2H, BH<sub>t</sub>), 1.78 (s, 15H, C<sub>5</sub>Me<sub>5</sub>), 1.77 (s, 15H, C<sub>5</sub>Me<sub>5</sub>) ppm. <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, C<sub>6</sub>D<sub>6</sub>): 103.7 (s, C<sub>5</sub>Me<sub>5</sub>), 102.0 (s, C<sub>5</sub>Me<sub>5</sub>), 9.8 (s, C<sub>5</sub>Me<sub>5</sub>), 9.1 (s, C<sub>5</sub>Me<sub>5</sub>) ppm. FTIR (CH<sub>2</sub>Cl<sub>2</sub>): 2461 (BH<sub>t</sub>), 1996 (CO) cm<sup>-1</sup>.

**Synthesis of 3:** In a flame-dried Schlenk tube, *nido-II* (0.05 g, 0.096 mmol) was stirred with [BF<sub>3</sub>.OEt<sub>2</sub>] (0.1 mL, 0.2 mmol) in THF at room temperature for 1 h. The volatile components were removed under vacuum. The residue was extracted into hexane:CH<sub>2</sub>Cl<sub>2</sub> mixture after removal of solvent and passed through a frit using celite. The residue was



subjected to chromatographic purification using silica gel TLC plates. Elution with a mixture of hexane:dichloromethane (40:60 v/v) in inert atmosphere yielded blue **3** (0.04 g, 46 %)

**3:** MS (ESI<sup>+</sup>):  $m/z$  calculated for  $[C_{30}H_{46}B_2O_2F_6Ru_3+Na]^+$ : 902.0654, found: 902.0642. <sup>19</sup>F NMR (470 MHz, C<sub>6</sub>D<sub>6</sub>, 22 °C):  $\delta$  = -146.16 (br, <sup>10</sup>BF<sub>3</sub>), -146.22 (br, <sup>11</sup>BF<sub>3</sub>) ppm. <sup>11</sup>B{<sup>1</sup>H} NMR (160 MHz, C<sub>6</sub>D<sub>6</sub>, 22 °C):  $\delta$  = 1.0 (s, 2B) ppm. <sup>1</sup>H NMR (500 MHz, C<sub>6</sub>D<sub>6</sub>, 22 °C):  $\delta$  = 1.63 (s, 30H, C<sub>5</sub>Me<sub>5</sub>), 1.62 (s, 15H, C<sub>5</sub>Me<sub>5</sub>), -15.68 (s, 1H, Ru-H-Ru) ppm. <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>, 22 °C):  $\delta$  = 95.2 (s, C<sub>5</sub>Me<sub>5</sub>), 93.4, (s, C<sub>5</sub>Me<sub>5</sub>), 11.1 (s, C<sub>5</sub>Me<sub>5</sub>), 11.0 (s, C<sub>5</sub>Me<sub>5</sub>) ppm. FTIR (CH<sub>2</sub>Cl<sub>2</sub>): 1433, 1377 (B-O), 1266, 882 (B-F) cm<sup>-1</sup>.

#### X-ray structure Determination.

Crystallographic information of compounds **1-3** is given below. Suitable crystals of **1**, **2** and **3** were produced by slow diffusion of dichloromethane-hexane mixture. The crystal data of **1** was collected and integrated by using a D8 VENTURE

Bruker AXS diffractometer at 150(2) K, **2** using Bruker axs Kappa Apex-II CCD diffractometer at 296(2) K and **3** using Bruker axs kappa Apex3 CMOS diffractometer at 299(2) K with graphite monochromated Mo-K $\alpha$  ( $\lambda$  = 0.71073 Å) radiation. The structures were solved by dual-space algorithm using SHELXT program<sup>28</sup> and refined with full-matrix least-squares methods based on F<sup>2</sup> (SHELXL program). Olex2 software used for the drawing of all the compounds.<sup>29</sup>

Crystal data for **1**: CCDC 2048248, C<sub>26</sub>H<sub>34</sub>B<sub>2</sub>FeO<sub>6</sub>W<sub>2</sub>, Mr = 887.70, monoclinic,  $P2_1/c$ ,  $a$  = 16.3632(19) Å,  $b$  = 9.9059(11) Å,  $c$  = 17.6686(18) Å,  $\alpha$  = 90°,  $\beta$  = 91.271(4)°,  $\gamma$  = 90°,  $V$  = 2863.2(5) Å<sup>3</sup>,  $Z$  = 4,  $\rho_{calcd}$  = 2.059 mg/m<sup>3</sup>,  $\mu$  = 8.552 mm<sup>-1</sup>,  $F(000)$  = 1688,  $R1$  = 0.0359,  $wR2$  = 0.0965, 6549 independent reflections [ $2\theta \leq 55.0^\circ$ ] and 357 parameters.

Crystal data for **2**: CCDC 2052992, C<sub>26</sub>H<sub>32</sub>B<sub>2</sub>Co<sub>2</sub>O<sub>6</sub>Rh<sub>2</sub>, Mr = 785.81, monoclinic,  $P2_1/n$ ,  $a$  = 11.319(5) Å,  $b$  = 15.288(7) Å,  $c$  = 16.746(7) Å,  $\alpha$  = 90°,  $\beta$  = 93.575(17)°,  $\gamma$  = 90°,  $V$  =





2892(2) Å<sup>3</sup>,  $Z = 4$ ,  $\rho_{\text{calcd}} = 1.805$  mg/m<sup>3</sup>,  $\mu = 2.286$  mm<sup>-1</sup>,  $F(000) = 1560$ ,  $R1 = 0.1152$ ,  $wR2 = 0.2842$ , 4825 independent reflections [ $2\theta \leq 49.98^\circ$ ] and 354 parameters.

Crystal data for **3**: CCDC 2045991, C<sub>30</sub>H<sub>45</sub>B<sub>2</sub>F<sub>6</sub>O<sub>2</sub>Ru<sub>3</sub>,  $M_r = 876.49$ , Orthorhombic, *Aba2*,  $a = 18.1307(11)$  Å,  $b = 34.026(2)$  Å,  $c = 21.9349(13)$  Å,  $\alpha = 90^\circ$ ,  $\beta = 90^\circ$ ,  $\gamma = 90^\circ$ ,  $V = 13532.0(14)$  Å<sup>3</sup>,  $Z = 16$ ,  $\rho_{\text{calcd}} = 1.721$  mg/m<sup>3</sup>,  $\mu = 1.381$  mm<sup>-1</sup>,  $F(000) = 6992.908$ ,  $R1 = 0.0395$ ,  $wR2 = 0.0891$ , 11883 independent reflections [ $2\theta \leq 50.0^\circ$ ] and 1005 parameters.

## Results and discussion

*Bis-{hydrido(borylene)} cluster* [ $(\mu_3\text{BH})_2\text{H}_2\{\text{Cp}^*\text{W}(\text{CO})_2\}\{\text{Cp}^*\text{W}(\text{CO})\}\{\text{Fe}(\text{CO})_3\}$ ] (**1**):

As shown in Scheme 2, thermolysis of an *in situ* obtained intermediate, generated from the


reaction of  $[\text{Cp}^*\text{WCl}_4]$  and  $[\text{LiBH}_4\text{THF}]$ , with  $\text{Fe}_2(\text{CO})_9$  yielded yellow solid **1** along with the earlier reported clusters

$[\{\text{Cp}^*\text{W}(\text{CO})_2\}_2\text{H}_2(\mu_3\text{-BH})_2\{\text{Fe}(\text{CO})_2\}]$ ,<sup>26</sup>

$[\{\text{Cp}^*\text{W}(\text{CO})_2\}_2(\mu_3\text{-BH})_2\{\text{Fe}(\text{CO})_3\}]$ ,<sup>26</sup>

and  $[(\text{Cp}^*\text{W})_2\text{B}_5\text{H}_9]$ .<sup>27</sup> The ESI-Mass

spectrum of **1** revealed a molecular ion peak at  $m/z$  829.0269. The  $^{11}\text{B}\{^1\text{H}\}$  NMR spectrum of compound **1** shows signals at  $\delta = 87.9$  and 62.4 ppm in 1:1 ratio, signifying two equivalent of boron atoms. The  $^1\text{H}$  NMR spectrum displays two signals at  $\delta = 2.19$  and 1.97 ppm indicate two different types of Cp\* ligands. Besides signals due to two B-H terminal protons, the  $^1\text{H}$  NMR shows two upfield chemical shifts at  $\delta = -10.64$  and  $-12.44$  ppm which demonstrates the existence of bridging W-H-B protons.




**Scheme 2.** Synthesis of compound **1**.

The existence of Cp\* ligands has also been confirmed by  $^{13}\text{C}\{^1\text{H}\}$  NMR. The infrared spectra (IR) of **1** shows the stretching frequency for terminal B-H ( $2415\text{ cm}^{-1}$ ) and carbonyl groups ( $2014, 1943\text{ cm}^{-1}$ ).

In order to determine the solid-state structure of **1**, single-crystal X-ray analysis was performed on a suitable crystal obtained from a solution of compound **1** in  $\text{CH}_2\text{Cl}_2$  and hexane solution at  $-2^\circ\text{C}$ . The X-ray structure of **1**, shown in Fig. 1. with the molecular formula as  $[\text{H}_2(\mu_3\text{-BH})_2\{\text{Cp}^*\text{W}(\text{CO})_2\}\{\text{Cp}^*\text{W}(\text{CO})\}\{\text{Fe}(\text{CO})_3\}]$  signifies a trigonal bipyramid  $\text{W}_2\text{FeB}_2$  core, in which one Fe and two

W atoms occupied the equatorial positions and two boron atoms are in the axial positions. The W-W bond distance of  $2.9946(4)\text{ \AA}$  is marginally longer in **1** compared to W-W bond distance of reported ditungstaboranes clusters.<sup>26-27,30</sup> The average W-Fe and Fe-B bond distances of **1**  $2.758(9)$  and  $2.079(7)$  respectively, are consistent with the W-Fe and Fe-B bond distances ( $2.77(11)$  and  $2.04(9)$ ) of  $[\{\text{Cp}^*\text{W}(\text{CO})_2\}_2\text{H}_2(\mu_3\text{-BH})_2\{\text{Fe}(\text{CO})_2\}]$ <sup>26</sup>. Compound **1** has 6 skeleton electron pairs (SEP) count which is in accord with the *closo*-trigonal bipyramidal (tbp) geometry.<sup>31</sup>



**Fig. 1.** Molecular structure and labelling diagram of **1**. Selected bond lengths ( $\text{\AA}$ ) and angles ( $^\circ$ ): W1-W2  $2.9946(4)$ , Fe1-W1  $2.7592(9)$ , Fe1-W2  $2.7577(9)$ , W2-B51




2.325(7), W2-B52 2.298(8), B51-W1 2.332(7), B52-W1 2.340(6), Fe1-B52 2.054(7), Fe1-B51 2.105(7), W1-B52-Fe1 77.5(2), W2-B51-Fe1 76.8(2).

On the basis of available literature, cluster **1** can be considered as *bis*-{hydrido(borylene)} species. Braunschweig group reported a new class of mono- and dinuclear borylene compound called hydridoborylene by intermetallic borylene transfer reaction in 2013.<sup>32</sup> In 2017, Tobita *et.al.* synthesized base-stabilized Hydrido (hydroborylene) tungsten complexes where W-B bond is bridged by a hydrido ligand.<sup>33</sup> The core geometry of **1** is analogous to that of *bis*{hydrido(borylene)} species  $[H_2(\mu_3-BH)_2\{Cp^*W(CO)_2\}_2\{Fe(CO)_2\}]$  recently reported by us.<sup>26</sup> In cluster **1**, the W2 atom contains one CO ligand, and W1 atom contains two CO ligands. But in the case of reported *bis*-{hydrido(borylene)} species both tungsten atom comprises two CO ligands. One of the unique features of

compound **1** is both the hydrido ligands are attached to one tungsten (W2) atom. Compound **1** is the rare example of transition metal borylene complex containing a triply-bridging *bis*-{hydrido(borylene)} ligand.

*Reactivity of nido-[Cp\*Rh<sub>2</sub>(B<sub>3</sub>H<sub>7</sub>)] with Co<sub>2</sub>(CO)<sub>8</sub>:*

Further to synthesized metal-rich rhodaborane we explored the reaction of *nido*-[Cp\*Rh<sub>2</sub>(B<sub>3</sub>H<sub>7</sub>)] (**I**) with Co<sub>2</sub>(CO)<sub>8</sub> at room temperature that isolates black compound **2** in moderate yield. The mass spectrum of **2** shows a molecular ion peak at *m/z* 808.9052. The <sup>11</sup>B{<sup>1</sup>H} NMR spectrum displayed two resonances at  $\delta = 168.0$  and  $152.5$  ppm. The <sup>1</sup>H NMR spectrum discloses the existence of two Cp\* ligands at  $\delta = 1.78$  and  $1.77$  ppm and signals for B-H terminal protons at  $\delta = 9.35$  ppm.




**Scheme 3.** Synthesis of compound **2**.

The IR spectrum of **2** features the stretching frequency for terminal B-H ( $2461\text{ cm}^{-1}$ ) and carbonyl groups ( $1996\text{ cm}^{-1}$ ).

As we were unable to get exact structure from these spectroscopic data,

X-ray diffraction analysis was undertaken to determine the solid-state structure on an appropriate crystal of compound **2** obtained at -2 °C. in hexane and CH<sub>2</sub>Cl<sub>2</sub> solution.






**Fig. 2.** Molecular structure and labelling diagram of **2**. Selected bond lengths (Å) and angles (°): Rh1-Rh2 2.689(2), B1-B2 1.84(3), B1-Rh1 2.07(2), B1-Co1 2.07(2), B1-Rh2 2.08(2), B1-H1 1.1000, B2-Co1 2.03(2), B2-Co2 2.08(2), B2-Rh2 2.11(3); B2-B1-Rh1 103.7(12), B1-B2-Co2 95.6(13), B2-Co2-Rh1 83.1(7), B1-Rh1-Co2 77.6(6), Co1-B1-Rh2 116.0(10), Co1-B2-Rh2 116.3(12), Co1-Rh1-Rh2 82.50(8).

The crystal structure of **2**, shown in Fig. 2, signifies an octahedral geometry consisting  $\text{Rh}_2\text{Co}_2\text{B}_2$ , in which one rhodium atom, Rh1, two boron atoms, B1 and B2 and one cobalt atom Co2, represents the equatorial plane and while Rh2 and Co1 atoms occupy the apical position. One carbonyl capping the triangular Rh-Co-Rh face. The Rh-Rh and Rh-B bond distances in compound **2** are consistent compared to Rh-Rh and Rh-B bond distances reported rhodaboranes.<sup>9b,16a,22a,26</sup> *Nido-I* having (n+2) SEP goes on for the formation of a *closo*-rhodaborane with (n+1) SEP.<sup>31</sup> Compound **2** has 7 skeleton electron pairs (SEP) count which is in accord with the *closo*-octahedral geometry. Compound **2** is interesting. It's a nice skeletal isomer of *closo*- $\text{M}_4\text{E}_2$  clusters discussed in the past (M= Ru or Os, E= CR or Bi).<sup>34</sup>

#### Synthesis of oxoruthena cluster **3**:

In an attempt to isolate metal-rich ruthenaboranes, we carried out the reaction of *nido*- $[\text{B}_3\text{H}_9(\text{Cp}^*\text{Ru})_2]$  (**II**) with mono-borane reagent  $[\text{BH}_3 \cdot \text{THF}]$  at 60 °C in THF for 15 h. This leads to the formation of known cluster *nido*- $[1,2-(\text{Cp}^*\text{Ru})_2\text{B}_4\text{H}_9(\mu\text{-H})]$ .<sup>24</sup> Besides, we have isolated a blue compound **3** in 46% yield.




**Scheme 4:** Synthesis of compound **3**.


This compound has been characterized by IR spectroscopy, Mass spectrometry, multi-nuclear NMR spectroscopy, and single-crystal X-ray diffraction analyses. Compound **3** exhibits a sharp peak at  $\delta = 1.0$  ppm in  $^{11}\text{B}\{^1\text{H}\}$  NMR spectrum. The  $^1\text{H}$  NMR



indicates the presence of two types of Cp\* ligands in 1:2 ratio at  $\delta = 1.62$  and 1.63 ppm respectively and an upfield



resonance at  $\delta = -15.68$  ppm due to the presence of Ru-H-Ru. Further the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum of **3** confirms the existence of two types of Cp\* ligands. Based upon the spectroscopic data, we can't conclude the structure. The structure of **3** became clear from the single crystal x-ray diffraction which determined as a trinuclear bis-( $\mu_3$ -oxo) ruthena cluster  $[(\text{Cp}^*\text{Ru})_3(\mu_3\text{-OBF}_3)_2(\mu\text{-H})]$ .



**Fig. 3.** Molecular structure and labelling diagram of **3** (One of the Cp\* ligand attached to Ru1 is omitted for clarity). Selected bond lengths ( $\text{\AA}$ ) and angles ( $^\circ$ ): Ru1-Ru2 2.834(2), Ru1-Ru3 2.837(17), Ru2-Ru3 2.822(2), Ru1-O1 2.024(9), Ru2-O2 2.079(10), Ru3-O2 2.034(10), O1-B1 1.531(12), O2-B2 1.514(14), Ru1-Ru2-Ru3 60.22(5), Ru3-O2-Ru1 88.3(5).



The core structure of **3** is similar to that of  $[(\text{Cp}^*\text{Ru})_3(\mu_3\text{-O})_2(\mu\text{-H})]^{35}$  and  $[(\text{CpRu})_3(\mu_3\text{-O})_2(\mu\text{-Cl})]^{36}$ . As shown in Fig. 3, each oxygen atom is connected to three Ru metals and one  $\text{BF}_3$  fragment. The molecule has a mirror plane passing through Ru3, O1, O2, B1, B2 and center of Ru1–Ru2 bond. Compound **3** can be considered as cluster having trigonal bipyramidal

geometry with *exo*- $\text{BF}_3$  fragment. The total electron count (TEC) of compound **3** is 48 and SEP count is 6. To find out the source of O- $\text{BF}_3$  fragment, we explore the reaction of *nido*- $[(\text{Cp}^*\text{Ru})_2\text{B}_3\text{H}_9]$  with  $\text{BF}_3\cdot\text{OEt}_2$ . After adding  $\text{BF}_3\cdot\text{OEt}_2$  to *nido*- $[(\text{Cp}^*\text{Ru})_2\text{B}_3\text{H}_9]$  in THF, the reaction mixture was stirred for 1 h at room temperature.

**Fig. 4.** (a)  $^{11}\text{B}\{^1\text{H}\}$  NMR spectrum (b)  $^{19}\text{F}$  NMR spectrum of **3** in  $\text{C}_6\text{D}_6$ .

A visible color change from orange to dark brown solution was observed upon the formation of **3**. The  $^{19}\text{F}$  NMR spectrum shows two resonances, one at  $\delta = -146.16$  ppm for the coupling with  $^{10}\text{B}$  nuclei and another resonance at  $\delta = -146.22$  ppm due to the coupling with  $^{11}\text{B}$  nuclei ( $J(^{11}\text{B}\text{-}^{19}\text{F}) = 6.2$  Hz) (Fig. 4).<sup>37</sup> The mass spectrometric analysis shows a molecular ion peak at  $m/z$  902.0642 which corresponds to the molecular formula  $[\text{C}_{30}\text{H}_{46}\text{B}_2\text{O}_2\text{F}_6\text{Ru}_3+\text{Na}]^+$ . The infrared spectrum indicates the presence of B-O stretching frequencies at 1377, 1433  $\text{cm}^{-1}$ . The average B–O bond distance of 1.522 Å is comparable to the

reported metallaoxaborane clusters.<sup>38</sup> The average Ru–Ru bond distance (2.831 Å) of **3** is slightly longer than  $[(\text{Cp}^*\text{Ru})_3(\mu_3\text{-O})_2(\mu\text{-H})]^{36}$  (2.7135(10) Å). In compound **3** the average Ru–O distances of 2.045 Å and 2.050 Å for Ru–O1 and Ru–O2 respectively are significantly longer than the bis( $\mu_3$ -oxo) complex  $[(\text{Cp}^*\text{Ru})_3(\mu_3\text{-O})_2(\mu\text{-H})]$  (2.004(6) Å). These bond length values lie within the range of metal-oxygen bond in reported bis( $\mu_3$ -oxo) complexes of transition metals.

## Conclusions

In this article, we have described the synthesis and characterization of



J. Indian Chem. Soc.

ISSN: 0019-4522

Vol..../Issue.../pp...



several metal-rich metallaboranes based on the reaction of cyclopentadienyl metal chlorides with boranes and metal carbonyls. Compound **1** represents one of the rare examples of transition metal borylene cluster containing a triply-bridging *bis*-{hydrido(borylene)} ligand. An *closo*-octahedral rhodaborane was synthesized from the reaction of *nido*-**I** with  $\text{Co}_2(\text{CO})_8$ . Also, a trinuclear  $\text{bis}(\mu_3\text{-oxo})$  metalla cluster having trigonal bipyramidal geometry with *exo*- $\text{BF}_3$  fragment was synthesized. All the synthesized clusters obey Wade-Mingos rule of the polyhedral skeleton electron count and are unique considering their cluster geometries.<sup>31</sup>

### Conflict of interest

The authors declare no conflict of interest.

### Acknowledgments

Generous support of the Council of Scientific & Industrial Research (Project No. 01(2939)/18/emr-ii), New Delhi, India, is gratefully acknowledged. R. B. and S. G. thank

IIT Madras for fellowship, A. N. P. thank UGC, India for fellowship.

### References

- (a) T. P. Fehlner, J.-F. Halet and J.-Y. Saillard, *Molecular Clusters. A Bridge to Solid-State Chemistry*, Cambridge University Press, Cambridge, 2007; (b) R. N. Grimes, *Metal Interactions with Boron Clusters*, Plenum Press, New York, 1982; (c) R. Borthakur, K. Saha, S. Kar and S. Ghosh, *Coord. Chem. Rev.*, 2019, **399**, 213021.
- (a) T. Onak, In: Wilkinson G, Abel E, Stone FGA (eds) *Comprehensive organometallic chemistry II*, chap. 6, vol 1. Pergamon Press, Oxford, p 217, 1995; (b) S. Kar, A. N. Pradhan and S. Ghosh, *Coord. Chem. Rev.*, 2021, **436**, 213796.
- N. S. Hosmane In: Siebert W (ed) *Advances in boron chemistry*. Royal Society of Chemistry, Cambridge, p 349, 1997.
- (a) J. L. Little, G. D. Friesen and L. J. Todd, *Inorg. Chem.*, 1977, **16**, 869; (b) R. P. Micciche, J. J. Briguglio and L. G. Sneddon, *Inorg. Chem.*, 1984, **23**, 3992.
- (a) S. Ghosh, B. C. Noll and T. P. Fehlner, *Dalton Trans.*, 2008, 371; (b) D. K. Roy, S. Ghosh and J.-F.





- Halet, *J. Cluster. Sci.*, 2014, **25**, 225.
6. (a) T. P. Fehlner, *Organometallics*, 2000, **19**, 2643; (b) S. Ghosh, A. M. Beatty and T. P. Fehlner, *Collect. Czech Chem. Commun.*, 2002, **67**, 808; (c) S. Ghosh, B. C. Noll and T. P. Fehlner, *Angew. Chem. Int. Ed.*, 2005, **44**, 6568; (d) D. K. Roy, S. K. Bose, R. S. Anju, V. Ramkumar and S. Ghosh, *Inorg. Chem.*, 2012, **51**, 10715; (e) S. K. Bose, K. Geetharani, B. Varghese and S. Ghosh, *Inorg. Chem.*, 2010, **49**, 6375.
  7. (a) S. Ghosh, X. Lei, M. Shang and T. P. Fehlner, *Inorg. Chem.*, 2000, **39**, 5373; (b) S. K. Bose, K. Geetharani, V. Ramkumar, S. M. Mobin and S. Ghosh, *Chem. Eur J.*, 2009, **15**, 13483; (c) R. S. Dhayal, S. Sahoo, K. H. K. Reddy, S. M. Mobin, E. D. Jemmis and S. Ghosh, *Inorg. Chem.*, 2010, **49**, 900; (d) S. Sahoo, R. S. Dhayal, B. Varghese and S. Ghosh, *Organometallics*, 2009, **28**, 1586.
  8. (a) T. P. Fehlner, *Proc. Indian Natl. Sci. Acad.*, 2002, **68**, 579; (b) S. Ghosh, M. Shang and T. P. Fehlner, *J. Am. Chem. Soc.*, 1999, **121**, 7451; (c) R. S. Anju, K. Saha, B. Mondal, V. Dorcet, T. Roisnel, J. - F. Halet, and S. Ghosh, *Inorg. Chem.*, 2014, **53**, 10527.
  9. (a) D. K. Roy, S. K. Bose, R. S. Anju, B. Mondal, V. Ramkumar and S. Ghosh, *Angew. Chem. Int. Ed.*, 2013, **52**, 3222; (b) D. K. Roy, R. Borthakur, R. Prakash, S. Bhattacharya, R. Jagan and S. Ghosh, *Inorg. Chem.*, 2016, **55**, 4764.
  10. (a) S. Ghosh, A. M. Beatty and T. P. Fehlner, *Angew. Chem. Int. Ed.*, 2003, **42**, 4678; (b) B. Le Guennic, H. Jiao, S. Kahlal, J.-Y. Saillard, J.-F. Halet, S. Ghosh, A. M. Beatty, A. L. Rheingold and T. P. Fehlner, *J. Am. Chem. Soc.*, 2004, **126**, 3203.
  11. (a) K. Geetharani, S. K. Bose, S. Sahoo, B. Varghese, S. M. Mobin and S. Ghosh, *Inorg. Chem.*, 2011, **50**, 5824; (b) R. Bag, B. Mondal, K. Bakthavachalam, T. Roisnel and S. Ghosh *Pure Appl. Chem.* 2018, **90**, 665.
  12. (a) N. N. Greenwood, A. Earnshaw, *Chemistry of the Elements*, 2nd ed., Butterworth- Heinemann, Oxford, 1997; (b) J. D. Kennedy, *Prog. Inorg. Chem.*, 1984, **32**, 519; (c) J. D. Kennedy, *Prog. Inorg. Chem.* 1986, **34**, 211.
  13. (a) A. S. Weller, "d and f-Block Metallaboranes" in *Comprehensive Organometallic Chemistry III* (Eds.: R. H. Crabtree, D. M. P. Mingos), Elsevier, Oxford, vol. 3,



- pp. 133, 2006; (b) N. S. Hosmane and J. A. Maguire in *Comprehensive Organometallic Chemistry III* (Eds.: R. H. Crabtree, D. M. P. Mingos), Elsevier, Oxford, vol. 3, pp. 175, 2007.
14. (a) Z. Yinghuai and N. S. Hosmane, *J. Organomet. Chem.*, 2013, **747**, 25; (b) N. S. Hosmane, J. A. Maguire and Z. Yinghuai, *Main Group Chem.*, 2006, **5**, 251; (c) S. Ghosh, A. L. Rheingold and T. P. Fehlner, *Chem. Commun.*, 2001, 895.
  15. (a) C. E. Housecroft, *Coord. Chem. Rev.*, 1995, **143**, 297; (b) K. Yuvaraj, D. K. Roy, K. Geetharani, B. Mondal, V. P. Anju, P. Shankhari, V. Ramkumar and S. Ghosh, *Organometallics*, 2013, **32**, 2705; (c) D. Y. Kim and G. S. Girolami, *J. Am. Chem. Soc.*, 2006, **128**, 10969; (d) J. Feilong and T. P. Fehlner *J. Am. Chem. Soc.*, 1987, **109**, 1860; (e) R. Okamura, K. Tada, K. Matsubara, M. Oshima and H. Suzuki, *Organometallics*, 2001, **20**, 4772.
  16. (a) K. Yuvaraj, M. Bhattacharyya, R. Prakash, V. Ramkumar and S. Ghosh, *Chem. Eur. J.*, 2016, **22**, 8889; (b) J. R. Pipal and R. N. Grimes, *Inorg. Chem.*, 1979, **18**, 252; (c) X. Lei, M. Shang and T. P. Fehlner, *Organometallics*, 2000, **19**, 4429; (d) K. Geetharani, S. K. Bose, B. Varghese and S. Ghosh, *Chem. Eur. J.*, 2010, **16**, 11357; (e) S. Gomosta, S. Kar, A. N. Pradhan, S. Bairagi, V. Ramkumar, and S. Ghosh, *Organometallics*, 2021, **40**, 529.
  17. (a) Z. Wang, H. Ye, Y. Li, Y. Li and H. Yan, *J. Am. Chem. Soc.*, 2013, **135**, 11289; (b) L. Deng, H.-S. Chan and Z. Xie, *Angew. Chem. Int. Ed.*, 2005, **44**, 2128; (c) S. K. Bose, D. K. Roy, P. Shankhari, K. Yuvaraj, B. Mondal, A. Sikder and S. Ghosh, *Chem. Eur. J.*, 2013, **19**, 2337.
  18. (a) J. Zhang and Z. Xie, *Acc. Chem. Res.*, 2014, **47**, 1623; (b) R. D. McIntosh, D. Ellis, G. M. Rosair and A. J. Welch, *Angew. Chem. Int. Ed.*, 2006, **45**, 4313; (c) A. Thakur, S. Sao, V. Ramkumar, and S. Ghosh, *Inorg. Chem.*, 2012, **51**, 8322.
  19. (a) J. B. Levine, S. H. Tolbert and R. B. Kaner, *Adv. Funct. Mater.*, 2009, **19**, 3519; (b) A. Friedrich, B. Winkler, E. A. Juarez-Arellano and L. Bayarjargal, *Materials*, 2011, **4**, 1648.
  20. (a) J. M. Vandenberg, B. T. Matthias, E. Corenzwit and H. Barz, *Mater. Res. Bull.*, 1975, **10**, 889; (b) K. Togano, P. Badica, Y. Nakamori, S. Orimo, H. Takeya



- and K. Hirata, *Phys. Rev. Lett.*, 2004, **93**, 247004.
21. B. P. T. Fokwa, *Eur. J. Inorg. Chem.*, 2010, 3075.
  22. (a) D. K. Roy, S. K. Barik, B. Mondal, B. Varghese and S. Ghosh, *Inorg. Chem.*, 2014, **53**, 667; (b) D. Sharmila, K. Yuvaraj, S. K. Barik, D. K. Roy, K. K. Chakrahari, R. Ramalakshmi, B. Mondal, B. Varghese and S. Ghosh, *Chem. Eur. J.*, 2013, **19**, 15219.
  23. M. L. H. Green, J. D. Hubert and P. J. Mountford, *Chem. Soc., Dalton Trans.*, 1990, 3793.
  24. X. Lei, M. Shang and T. P. Fehlner, *J. Am. Chem. Soc.*, 1999, **121**, 1275.
  25. G. E. Ryschkewitsch and K. C. Nainan, *Inorg. Synth.*, 1974, **15**, 113.
  26. R. Bag, S. Kar, S. Saha, S. Gomosta, B. Raghavendra, T. Roisnel and S. Ghosh, *Chem. Asian J.*, 2020, **15**, 780.
  27. A. S. Weller, M. Shang and T. P. Fehlner, *Organometallics*, 1999, **18**, 53.
  28. G. M. Sheldrick, *Acta Cryst.*, 2015, **A71**, 3.
  29. O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Cryst.*, 2009, **42**, 339.
  30. (a) B. Mandal, R. Bag, T. Rosinel and S. Ghosh *Inorg. Chem.*, 2019, **58**, 2744; (b) R. Bag, S. Saha, R. Borthakur, B. Mondal, T. Roisnel, V. Dorcet, J.-F. Halet and S. Ghosh, *Inorganics*, 2019, **7**, 27.
  31. (a) K. Wade, *J. Chem. Soc. D*, 1971, 792; (b) K. Wade, *Adv. Inorg. Chem. Radiochem.*, 1976, **18**, 1; (c) M. A. Fox and K. Wade, *Pure Appl. Chem.*, 2003, **75**, 1315.
  32. J. Bauer, S. Bertsch, H. Braunschweig, R. D. Dewhurst, K. Ferkinghoff, C. Hörl, K. Kraft and K. Radacki, *Chem. Eur. J.*, 2013, **19**, 17608.
  33. Z. Hui, T. Watanabe and H. Tobita, *Organometallics*, 2017, **36**, 4816.
  34. J. -F. Halet, *Coord. Chem. Rev.*, 1995, **143**, 637.
  35. H. Suzuki, T. Kakigano, K. Tada, M. Igarashi, K. Matsubara, A. Inagaki, M. Oshima, and T. Takao, *Bull. Chem. Soc. Jpn.*, 2005, **78**, 67.
  36. S. Takemoto, H. Ishii, M. Yamaguchi, A. Teramoto, M. Tsujita, D. Ozeki and H. Matsuzaka, *Organometallics*, 2019, **38**, 4298.
  37. E. L. Myers, C. P. Butts and V. K. Aggarwal, *Chem. Commun.*, 2006, 4434.
  38. (a) Y.-H. Kim, A. Brownless, P. A. Cooke, R. Greatrex, J. D. Kennedy

**J. Indian Chem. Soc.**

**ISSN: 0019-4522**

**Vol..../Issue.../pp...**



and M. Thornton-Pett, *Inorg. Chem. Commun.*, 1998, **1**, 19; (b) S. K. Bose, K. Geetharani, S. Sahoo, K. H. K. Reddy, B. Varghese, E. D. Jemmis and S. Ghosh, *Inorg. Chem.*, 2011, **50**, 9414.

J. Indian Chem. Soc.

ISSN: 0019-4522

Vol..../Issue.../pp...




---

## Table of Contents

### Synthesis and characterization of group 6-9 metal-rich homo- and hetero-metallaboranes

Ranjit Bag, Suman Gomosta, Alaka Nanda Pradhan, Thierry Roisnel and Sundargopal Ghosh\*



Metal rich metallaborane clusters with trigonal bipyramidal and octahedral geometry (see picture)

- Synthesis of metal-rich metallaboranes
- Rare examples of *bis*-{hydrido(borylene)}
- trigonal bipyramidal and octahedral geometry
- oxoruthena cluster

**Declaration of interests**

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: